

Determination of optimal grazing management for dairy cows in Galicia (Spain) using a decision support system.

O. P. Vázquez Yáñez, A. González Rodríguez and J. López Díaz

Centro de Investigaciones Agrarias de Mabegondo (CIAM), Apartado 10, 15080 A Coruña (Spain)

Email: orlando.vazquez.yanez@xunta.es

Keywords: dairy cows, decision support system, grazing management

Introduction GRAZEMORE is a decision support system (DSS) oriented to evaluate dairy grazing management decisions on pasture growth and milk production. The system integrates a herbage growth model (Barrett et al., 2003) and a herbage intake model (Delagarde et al., 2004) and was validated in several countries of the EU. The objective of this work was to use GRAZEMORE DSS to determine optimal grazing management of dairy cows at three supplementation levels and three grazing management strategies under the Galician conditions.

Materials and methods The GRAZEMORE DSS was used to simulate eight grazing scenarios using the average weather data of the CIAM in Galicia (NW Spain). The scenarios considered a herd of 50 dairy cows with a continuous calving pattern, grazing 13.6 ha divided into 17 paddocks from 1st May to 30th October. The potential peak milk yield was estimated to be 36 kg/d. The scenarios compared where three concentrate supplementation levels (0, 4 and 8 kg DM concentrate/d) and three management options (a daily pasture area offered (DPAO) of 0.2, 0.3 and 0.5 ha/d). The number of rotations during the grazing season was 7, 9 and 12 for a DPAO of 0.2, 0.3 and 0.5 ha/d respectively, but during the five initial rotations only half of paddocks were grazed and were assumed to be cut for silage.

Results Average pasture allowance, pasture intake, total intake, and daily milk production between 15th May to 30th October are shown in Table 1. These results show that increasing DPAO resulted in higher pasture allowance (from 15.7 to 36.3 kg DM/d) and, as a consequence, pasture intake was raised from 9.9 to 13.2 kg DM/d. Increasing supplementation level reduced pasture intake and the magnitude of the reduction was enlarged with increased pasture allowances (0.4 kg DM in low DPAO vs. 0.6 kg DM in high DPAO for a total supplementation of 4 kg DM/d). Daily milk production ranged from 9.8 kg/d for 0.2 ha/d DPAO and no concentrate, to 27.0 kg/d for 0.3 ha/d DPAO and 8 kg DM concentrate. Total silo production in low DPAO system (46 t DM) was almost twice the silo production in the high DPAO. Grass utilization (total pasture cut and consumed/pasture produced) increased with DPAO but decreased with concentrate supplementation level.

Table 1 Pasture and milk production for the scenarios simulated

Scenarios	Area offered (ha/d)							
	0.2			0.3			0.5	
Supplementation (Kg Conc/cow)	0	4	8	0	4	8	0	4
Pasture allowance (kg DM/cow/d)	15.7	16.3	17.1	21.3	22.5	24.4	33.3	36.3
Pasture DM intake (kg DM/cow/d)	9.9	9.5	8.8	11.4	10.9	9.8	13.2	12.6
Total DM intake (kg DM/cow/d)	9.9	13.5	16.8	11.4	14.9	17.8	13.2	16.6
Milk (kg/cow/d)	9.8	18.6	25.2	13.1	21.5	27.0	17.2	24.8
Total pasture intake (t DM)	94.9	91.0	83.6	108.7	104.1	93.7	126.0	120.5
Total silo cut (t DM)	46.4	46.4	46.5	33.2	33.2	33.3	23.9	23.9
Total grass produced (t DM)	165.9	166.1	166.3	161.4	161.7	162.0	160.4	160.8
Grass utilization (%)	85%	83%	78%	88%	85%	78%	93%	90%

Conclusions This work shows the sensibility of the grazing system to different grazing and supplementation strategies and the ability of GRAZEMORE to assist in optimal management decisions. Future work must consider the impact of the grazing management system in the economics of the farm considering the whole year.

References

- Barrett P.D., A.S. Laidlaw and C.S. Mayne (submitted 2003). An Herbage Growth Model to Predict Pasture Production in Predominant Perennial Ryegrass Swards for use in a Decision Support System to Aid Pasture Management in Northwest Europe. *The Journal of Agricultural Science*.
- Delagarde R., P. Faverdin, C. Baratte, M. Bailhache and J.L. Peyraud (2004). The Herbage Intake Model for Grazing Dairy Cows in the EU Grazemore Project. Proceedings of the 20th General Meeting of the European Grassland Federation, Luzern, Switzerland.